

12

EUROPEAN PATENT APPLICATION

21 Application number: **88201172.9**

22 Date of filing: **08.06.88**

51 Int. Cl.⁴: **C07D 401/14** , **C07D 401/12** ,
C07D 417/14 , **C07D 405/14** ,
C07D 409/14 , **C07D 471/04** ,
A61K 31/44 , **A61K 31/445** ,
A61K 31/415 , **A61K 31/425** ,
A61K 31/34

30 Priority: **19.06.87 US 63899**

43 Date of publication of application:
21.12.88 Bulletin 38/51


84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

71 Applicant: **JANSSEN PHARMACEUTICA N.V.**
Turnhoutseweg 30
B-2340 Beerse(BE)

72 Inventor: **Janssens, Frans Eduard**
Tinstraat 79
B-2830 Bonheiden(BE)
 Inventor: **Torremans, Joseph Leo Ghislanus**
Lijsterstraat 11
B-2340 Beerse(BE)
 Inventor: **Dieis, Gaston Stanislas Marcella**
Oosteinde 12
B-2380 Ravels(BE)

52 Novel **N-(4-piperidinyl)** bicyclic condensed 2-imidazolamine derivatives.

57 Novel **N-(4-piperidinyl)** bicyclic condensed 2-imidazolamine derivatives of formula



wherein: $A^1 = A^2 = A^3 = A^4$ is $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$, $-\text{N}=\text{CH}-\text{CH}=\text{CH}-$, $-\text{CH}=\text{N}-\text{CH}=\text{CH}-$, $-\text{CH}=\text{CH}-\text{N}=\text{CH}$ or $-\text{CH}=\text{CH}-\text{CH}=\text{N}-$; R is hydrogen or C_{1-6} alkyl; R^1 is furanyl substituted with C_{1-6} alkyl, pyrazinyl, thiazolyl, or imidazolyl optionally substituted with C_{1-6} alkyl; L is C_{3-6} alkenyl optionally substituted with Ar^3 , or L is a radical of formula $-\text{Alk}-\text{R}^3$, $-\text{Alk}-\text{O}-\text{R}^4$, $-\text{Alk}-\text{N}(\text{R}^5)-\text{R}^6$, $-\text{Alk}-\text{Z}-\text{C}(=\text{O})-\text{R}^7$ or $-\text{CH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{O}-\text{R}^9$; the pharmaceutically acceptable acid addition salts and possible stereochemically isomeric forms thereof, which compounds are anti-histaminic agents; pharmaceutical compositions containing such compounds as an active ingredient and methods of preparing said compounds and pharmaceutical compositions.


EP 0 295 742 A1

N vel **N**-(4-piperidinyl) bicyclic condensed 2-imidazolamine derivativesBackground of the invention

In the U.S. Patent No. 4,219,559 there are described a number of 1-substituted **N**-heterocycl-4-

piperidinamines as compounds having useful anti-histaminic properties.
Further series of **N**-heterocycl-4-piperidinamines as compounds having useful anti-histaminic and serotonin-antagonistic properties have been described in U.S. Patent Nos. 4,556,660, 4,634,704, 4,695,569 and 4,588,722.

The compounds of the present invention are substituted in a previously undisclosed manner and show

Description of the invention:

the pharmaceutically acceptable acid-addition salts and possible stereochemically isomeric forms thereof, wherein:

$A^1 = A^2 - A^3 = A^4$ is a bivalent radical having the formula

-CH=CH-CH=CH- (a-1),

-N=CH-CH=CH- (a-2),

-CH=N-CH=CH- (a-3),

-CH=CH-N=CH- (a-4), or

-CH=CH-CH=N- (a-5);

wherein one or two hydrogen atoms in said radicals (a-1)-(a-5) may, independently from each other, be replaced by halo, C_{1-6} alkyl, C_{1-6} alkyloxy, trifluoromethyl or hydroxy;

R is hydrogen or C_{1-6} alkyl;

R¹ is furanyl substituted with C_{1-6} alkyl, pyrazinyl, thiazolyl, or imidazolyl optionally substituted with C_{1-6} alkyl;

R² is hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl, Ar^3-C_{1-6} alkyl or $(C_{1-6}alkyl)-CO$;

L is C_{3-6} alkenyl optionally substituted with Ar^3 , or L is a radical of formula

-Alk-R³ (b-1);

-Alk-O-R⁴ (b-2),

-Alk-N(R⁵)-R⁶ (b-3),

-Alk-Z-C(=O)-R⁷ (b-4), or

-CH₂-CH(OH)-CH₂-O-R⁹ (b-5);

R³ is hydrogen, Ar^1 -thio, Ar^3 -sulfonyl, 4,5-dihydro-5-oxo-1H-tetrazol-1-yl being optionally substituted in its 4-position with Ar^3 or C_{1-6} alkyl, 2,3-dihydro-1,4-benzodioxin-2-yl, 2,3-dihydro-1,4-benzodioxin-6-yl, 4-morpholinyl, 1-piperidinyl or 1-pyrrolidinyl, or when R¹ is furanyl substituted with C_{1-6} alkyl and $A^1 = A^2 - A^3 = A^4$ is a bivalent radical of formula (a-1) or (a-2), R³ may also be Ar^1 , 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl or 1-($C_{1-6}alkyl$)pyrrolyl;

R⁴ is $C_{1-6}alkyl$ or Ar^1 ;

R⁵ is $C_{1-6}alkyl$ optionally substituted with Ar^2 ;

R⁶ is Ar^2 or $C_{1-6}alkyl$ optionally substituted with Ar^2 or when $A^1 = A^2 - A^3 = A^4$ is a radical of formula (a-1) or (a-2), R⁶ may also be 1H-benzimidazol-2-yl;

R⁷ is $C_{1-6}alkyl$, $Ar^2-C_{1-6}alkyl$, Ar^2 , amino, Ar^2 -amino, mono- and di($C_{1-6}alkyl$)amino, mono- and di($Ar^2-C_{1-6}alkyl$)amino, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl, $C_{3-6}cycloalkyloxy$, $C_{1-6}alkyloxy$ or Ar^1-C-

: -₅alkyloxy;

Z is O, NR⁸ or a direct bond; said R⁸ being hydrogen or C₁₋₅alkyl optionally substituted with Ar¹;

R⁹ is Ar³;

each Alk independently being C₁₋₅alkanediyl;

5 Ar¹ is phenyl, substituted phenyl, naphthalenyl, thienyl, halothienyl, C₁₋₅alkyl substituted thienyl or furanyl, wherein said substituted phenyl is phenyl substituted with 1, 2 or 3 substituents, each independently selected from the group consisting of halo, hydroxy, nitro, cyano, trifluoromethyl, C₁₋₅alkyl, C₁₋₅alkyloxy, C₁₋₆alkylthio, mercapto, C₁₋₆alkylsulfonyl, C₁₋₆alkylsulfonylC₁₋₆alkyl, phenylC₁₋₅alkylsulfonyl, phenylsulfonylC₁₋₅alkyl, amino, mono- and di(C₁₋₆alkyl)amino, carboxy, C₁₋₅alkyloxycarbonyl and C₁₋₆alkylcarbonyl;

10 Ar² has the same meanings of Ar¹, and may also be pyridinyl, mono- and di(C₁₋₅alkyloxy)pyridinyl or furanyl substituted with C₁₋₆alkyl; and

Ar³ has the same meanings of Ar², and may also be pyrazinyl, 1-(C₁₋₆alkyl)pyrrolyl, thiazolyl or imidazolyl optionally substituted with C₁₋₆alkyl;

15 provided that A¹ = A²-A³ = A⁴ is a radical of formula (a-1) or (a-2) when L is a radical of formula (b-4).

As used in the foregoing definitions the term halo is generic to fluoro, chloro, bromo and iodo; the term "C₁₋₆alkyl" is meant to include straight and branched saturated hydrocarbon radicals, having from 1 to 6 carbon atoms, such as, for example, methyl, ethyl, 1-methylethyl, 1,1-dimethylethyl, propyl, butyl, pentyl, 20 hexyl and the like; the term "C₃₋₆cycloalkyl" is generic to cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; the term "C₃₋₆alkenyl" is meant to include straight and branch chained hydrocarbon radicals containing one double bond and having from 3 to 6 carbon atoms such as, for example, 3-propenyl, 2-butenyl and the like, and when a C₃₋₆alkenyl is substituted on a heteroatom, then the carbon atom of said C₃₋₆alkenyl connected to said heteroatom preferably is saturated.

25 The compounds of formula (I) may also exist in hydrated or in solvent addition forms and said forms are intended to be included within the scope of the present invention.

An interesting subgroup among the compounds of formula (I) comprises those compounds of formula (I) wherein A¹ = A²-A³ = A⁴ is a bivalent radical having the formula (a-1). Another interesting subgroup among the compounds of formula (I) comprises those compounds of formula (I) wherein A¹ = A²-A³ = A⁴ is a bivalent 30 radical having a formula (a-2) through (a-5), with (a-2) being the most interesting subgroup.

Preferred compounds within the invention are those compounds of formula (I) wherein R is hydrogen and R² is hydrogen or C₁₋₆alkyl.

Particularly preferred compounds within the invention are those preferred compounds of formula (I) wherein L is C₃₋₆alkenyl optionally substituted with phenyl or substituted phenyl, or wherein L is a radical 35 of formula (b-1), (b-2), (b-4) or (b-5) wherein R³ is as defined hereinabove, R⁴ is C₁₋₆alkyl, phenyl or substituted phenyl, R⁷ is phenyl, substituted phenyl or C₁₋₆alkyl optionally substituted with phenyl or substituted phenyl, R⁸ is hydrogen or C₁₋₆alkyl, and R⁹ is phenyl, substituted phenyl or naphthalenyl.

More particularly preferred compounds within the invention are those particularly preferred compounds of formula (I) wherein L is a radical of formula (b-1) or (b-2) wherein R³ is hydrogen, phenylsulfonyl, 4,5- 40 dihydro-5-oxo-1H-tetrazol-1-yl being optionally substituted in its 4-position with C₁₋₄alkyl, 2,3-dihydro-1,4-benzodioxin-2-yl, 4-morpholinyl, 1-piperidinyl or 1-pyrrolidinyl, or when R¹ is furanyl substituted with C₁₋₆alkyl and A¹ = A²-A³ = A⁴ is a bivalent radical of formula (a-1) or (a-2), R³ may also be 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl, thienyl, furanyl, phenyl or substituted phenyl.

Especially preferred compounds within the invention are those more particularly preferred compounds 45 of formula (I) wherein A¹ = A²-A³ = A⁴ is a bivalent radical of formula (a-1) or (a-2) and L is a radical of formula (b-1) wherein R³ is hydrogen or phenyl optionally substituted with halo, C₁₋₄alkyl, C₁₋₄alkyloxy or hydroxy.


A particular subgroup of compounds of formula (I) comprises those compounds, preferred, particularly preferred, more particularly preferred or especially preferred compounds wherein R¹ is furanyl substituted 50 with C₁₋₆alkyl.

A more particular subgroup of compounds of formula (I) comprises those compounds, preferred, particularly preferred, more particularly preferred or especially preferred compounds of formula (I) wherein -Alk-R¹ is C₁₋₄alkyl-5-C₁₋₄alkyl-2-furanyl, C₁₋₄alkyl-4-C₁₋₄alkyl-2-furanyl, C₁₋₄alkyl-3-C₁₋₄alkyl-2-furanyl, C₁₋₄alkyl-2-C₁₋₄alkyl-3-furanyl, C₁₋₄alkyl-4-C₁₋₄alkyl-3-furanyl or C₁₋₄alkyl-5-C₁₋₄alkyl-3-furanyl.

55 The most preferred compounds within the invention are selected from the group consisting of 3-[(5-methyl-2-furanyl)methyl]-N-(1-methyl-4-piperidinyl)-3H-imidazo[4,5-b]pyridin-2-amine, the pharmaceutically acceptable acid addition salts and the stereochemically isomeric forms thereof.


In order to simplify the structural representations of the compounds of formula (I) and of certain

precursors and intermediates thereof the



radical will hereafter be represented by the symbol D.


The compounds of formula (I) are generally prepared by N-alkylating a piperidine of formula (II) with a reagent of formula (III).



In formula (III) and in a number of the following intermediates W and W' represent an appropriate leaving group such as, for example, halo, preferably, chloro, bromo or iodo, or a sulfonyloxy group such as, for example, methylsulfonyloxy or 4-methylphenylsulfonyloxy, whereas W may also be C₁₋₆alkyloxy and C₁₋₆alkylthio.

The above N-alkylation reaction is conveniently conducted in an inert organic solvent such as, for example, an aromatic hydrocarbon, e.g., benzene, methylbenzene, dimethylbenzene and the like; an alkanol, e.g., methanol, ethanol, 1-butanol and the like; a ketone, e.g., 2-propanone, 4-methyl-2-pentanone and the like; an ether, e.g., 1,4-dioxane, 1,1'-oxybisethane, tetrahydrofuran and the like; a polar aprotic solvent e.g., N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), dimethyl sulfoxide (DMSO), nitrobenzene, 1-methyl-2-pyrrolidinone, and the like. The addition of an appropriate base such as, for example, an alkali or an earth alkaline metal carbonate, hydrogen carbonate, hydroxide, amide or hydride, e.g., sodium carbonate, sodium hydrogen carbonate, potassium carbonate, sodium hydroxide, calcium carbonate, calcium hydroxide, sodium hydride, sodium amide and the like, or an organic base, such as, for example, a tertiary amine, e.g., N,N-diethylethanamine, N (1-methylethyl)-2-propanamine, 4-ethylmorpholine and the like may be utilized to pick up the acid which is liberated during the course of the reaction. In some instances the addition of an iodide salt, preferably an alkali metal iodide, is appropriate. Somewhat elevated temperatures may enhance the rate of the reaction.


The compounds of formula (I) wherein L is a radical of formula (b-1), (b-2), (b-3) or (b-4), said radical L being represented by the radical L'H-, and said compounds being represented by the formula (I-a), can also be prepared by the reductive N-alkylation reaction of (II) with an appropriate ketone or aldehyde of formula L¹=O (IV), said L¹=O being an intermediate of formula L¹H₂ wherein two geminal hydrogen atoms are replaced by =O; and L¹ is a geminal bivalent radical comprising R³-C₁₋₆alkylidene, R⁴-O-C₁₋₆alkylidene, R⁶R⁵-N-C₁₋₆alkylidene and R⁷-C(=O)-Z-C₁₋₆alkylidene.




Said reductive N-alkylation reaction may conveniently be carried out by catalytically hydrogenating a mixture of the reactants in a suitable reaction-inert organic solvent according to art-known catalytic hydrogenating procedures. The reaction mixture may be stirred and/or heated in order to enhance the reaction rate. Suitable solvents are, for example, water; lower alkanols, e.g. methanol, ethanol, 2-propanol and the like; cyclic ethers, e.g. 1,4-dioxane and the like; halogenated hydrocarbons, e.g. trichloromethane and the like; N,N-dimethylformamide; dimethyl sulfoxide and the like; or a mixture of two or more of such solvents. The term "art-known catalytic hydrogenating procedures" means that the reaction is carried out

under hydrogen atmosphere and in the presence of an appropriate catalyst such as, for example, palladium-on-charcoal, platinum-on-charcoal and the like. In order to prevent the undesired further hydrogenation of certain functional groups in the reactants and the reaction products it may be advantageous to add an appropriate catalyst-poison to the reaction mixture, e.g., thiophene and the like.

- 5 The compounds of formula (I) wherein L is a radical of formula (b-2) wherein R^4 is Ar^1 , said R^4 being represented by R^{4-a} and said compounds by formula (I-b), may also be prepared by O-alkylating an intermediate of formula (V) with a reagent of formula (VI).




- 15 The compounds of formula (I-b) can also be prepared by O alkylating an intermediate of formula (VII) with a reagent of formula (VIII).




The O-alkylation reactions of (VI) with (V) and (VIII) with (VII) may be carried out by stirring the reactants, preferably at somewhat elevated temperatures in an inert organic solvent such as, for example, an aromatic hydrocarbon, e.g., benzene, methylbenzene, dimethylbenzene; a ketone, e.g., 2-propanone, 4-methyl-2-pentanone; an ether, e.g., 1,4-dioxane, 1,1'-oxybisethane, tetrahydrofuran; and a polar aprotic solvent, e.g., N,N-dimethylformamide; N,N-dimethylacetamide; dimethyl sulfoxide; nitrobenzene; 1-methyl-2-pyrrolidinone; and the like. The addition of an appropriate base such as, for example, an alkali metal carbonate of hydrogen carbonate, sodium hydride or an organic base such as, for example, N,N-diethylethanamine or N-(1-methylethyl)-2-propanamine may be utilized to pick up the acid which is liberated during the course of the reaction.

35 The compounds of formula (I) wherein L is a radical of formula (b-4) wherein Z is NR^8 or O, said Z being represented by Z^1 and R^7 is amino, Ar^2 -amino, C_{1-6} alkyl-amino or Ar^2-C_{1-6} alkyl-amino, said R^7 being represented by $R^{7-a}-NH-$, and said compounds by formula (I-c-1), can be prepared by reacting an isocyanate of formula (X) with a reagent of formula (IX).



45 The compounds of formula (I), wherein L is a radical of formula (b-4) wherein Z is NH and R^7 is other than Ar^2 or C_{1-6} alkyl optionally substituted with Ar^2 , said R^7 being represented by R^{7-b} , and said compounds being represented by formula (I-c-2), can be prepared by reacting an isocyanate of formula (XI) with a reagent of formula (XII).




55 The reaction of (X) with (IX), or (XII) with (XI) is generally conducted in a suitable reaction-inert solvent such as, for example, an ether, e.g., tetrahydrofuran and the like. Elevated temperatures may be suitable to enhance the rate of the reaction.

The compounds of formula (I) wherein L is a radical of formula (b-4) wherein R⁷ is Ar² or C₁₋₆alkyl optionally substituted with Ar², and Z is NR⁸ or O, said R⁷ being represented by R^{7-c} and said compounds by formula (I-c-3), can be prepared by reacting a reagent of formula (XIII) or a functional derivative thereof with an intermediate of formula (IX).




The reaction of (XIII) with (IX) may generally be conducted following art-known esterification- or amidation reaction procedures. For example, the carboxylic acid may be converted into a reactive derivative, e.g., an anhydride or a carboxylic acid halide, which subsequently, is reacted with (IX); or by reacting (XIII) and (IX) with a suitable reagent capable of forming amides or esters, e.g., dicyclohexylcarbodiimide, 2-chloro-1-methylpyridinium iodide and the like. Said reactions are most conveniently conducted in a suitable solvent such as, for example, an ether, e.g., tetrahydrofuran, a halogenated hydrocarbon, e.g., dichloromethane, trichloromethane or a polar aprotic solvent. The addition of a base such as, N,N-diethylethanamine may be appropriate.

The compounds of formula (I) wherein L is a radical formula L²-C₂₋₆alkanediyl, said L² being Ar¹-thio, Ar³-sulfonyl, Ar², C₁₋₆alkylcarbonyl, Ar²-C₁₋₆alkylcarbonyl, Ar²-carbonyl, C₁₋₆alkyloxycarbonyl, Ar¹-C₁₋₆alkyloxycarbonyl or C₃₋₆cycloalkyloxycarbonyl, and said compounds being represented by formula (I-d), may be prepared by reacting an appropriate alkenylene of formula (XIV) with a piperidine of formula (II).




The compounds of formula (I) wherein L is a radical of formula (b-5), said compounds being represented by formula (I-e), may be prepared by reacting a reagent of formula (XV) with a piperidine of formula (II).



The reaction of (XIV) with (II), and (XV) with (II) may be conducted by stirring and, if desired, heating the reactants in a suitable solvent such as, for example, an alkanone, e.g., 2-propanone, 4-methyl-2-pentanone, an ether, e.g., tetrahydrofuran, 1,1'-oxybisethane, an alcohol, e.g., methanol, ethanol, 1-butanol, a polar aprotic solvent, e.g., N,N-dimethylformamide, N,N-dimethylacetamide and the like solvents.

Particular compounds wherein L is methyl, said compounds being represented by the formula (I-f), may also be prepared by reducing an intermediate of formula C₁₋₆alkyl-O-C(=O)-D (XVI) with an appropriate reductant such as, for example, lithium tetrahydroaluminate in a suitable reaction-inert organic solvent, such as a lower alkanol.


The compounds of formula (I) can also be prepared by N-alkylating an intermediate of formula (XVII) with an appropriate reagent of formula (XVIII).



10

The said N-alkylation is conveniently carried out according to the procedures described hereinabove for the preparation of (I) starting from (III) and (II).

The compounds of formula (I) may alternatively be prepared by the cyclodesulfurization reaction of an appropriate thiourea derivative of formula (XIX), which may in situ be formed by condensing an isothiocyanate of formula (XX) with a diamine of formula (XXI)




25

Said cyclodesulfurization reaction may be carried out by the reaction of (XIX) with an appropriate alkyl halide, preferably iodomethane in an appropriate reaction-inert organic solvent, e.g., a lower alkanol such as methanol, ethanol, 2-propanol and the like. Otherwise, the cyclodesulfurization reaction may be carried out by the reaction of (XIX) with an appropriate metal oxide or salt in an appropriate solvent according to art-known procedures. For example, the compounds of formula (I) can easily be prepared by the reaction of (XIX) with an appropriate Hg(II) or Pb(II) oxide or salt, such as, for example HgO, HgCl₂, Hg(OAc)₂, PbO or Pb(OAc)₂. In certain instances it may be appropriate to supplement the reaction mixture with a small amount of sulfur. Even so methanediimines, especially N,N'-methanetetraylbis[cyclohexanamine] may be used as cyclodesulfurizing agents.

35



Or, the compounds of formula (I) may be prepared by reacting a piperidine derivative of formula (XXII) with a benzimidazole derivative of formula (XXIII).



In (XXII) and (XXIII) Q¹ and Q² are selected so that during the reaction of (XXII) with (XXIII) the -NR²- moiety is formed connecting the piperidine and benzimidazole moiety.


For example Q¹ may be a radical -NHR² and Q² a radical -W or inversely Q¹ may be a radical -W' and Q² a radical -NHR².

55



The reactions of (XXII-a) with (XXIII-a) and of (XXII-b) with (XXIII-b) are conveniently conducted in an inert organic solvent such as, for example, an aromatic hydrocarbon, e.g., benzene, methylbenzene, dimethylbenzene; a lower alcohol, e.g., methanol, ethanol, 1-butanol; a ketone, e.g., 2-propanone, 4-methyl-2-pentanone; an ether, e.g., 1,4-dioxane, 1,1'-oxybisethane, tetrahydrofuran; N,N-dimethylformamide; N,N-dimethylacetamide; nitrobenzene; dimethyl sulfoxide; 1-methyl-2-pyrrolidinone; and the like. The addition of an appropriate base such as, for example, an alkali metal carbonate or hydrogen carbonate, sodium hydride or an organic base such as, for example, N,N-diethylethanamine or N-(1-methylethyl)-2-propanamine may be utilized to pick up the acid which is liberated during the course of the reaction. In some circumstances the addition of a iodide salt, preferably an alkali metal iodide, is appropriate. Somewhat elevated temperatures may enhance the rate of the reaction.

Or, Q¹ may be an oxo radical and Q² a radical -NHR².



The reaction of (XXII-x) with (XXIII-b) is conveniently carried out by stirring the reactants, preferably at somewhat elevated temperatures, in a suitable reaction-inert organic solvent with an appropriate reductant. Preferably, the piperidone of formula (XXII-c) is first reacted with the benzimidazoleamine of formula (XXIII-b) to form an enamine, which optionally may be isolated and further purified, and subsequently subjecting the said enamine to a reduction reaction. Suitable solvents are, for example, water; lower alcohols, e.g., methanol, ethanol, 2-propanol; cyclic ethers, e.g., 1,4-dioxane; halogenated hydrocarbons, e.g., trichloromethane; N,N-dimethylformamide; N,N-dimethylacetamide; dimethyl sulfoxide and the like; or a mixture of such solvents. Appropriate reductants are for example, metal or complex metal hydrides, e.g. sodium borohydride, lithium aluminiumhydride; or hydrogen, the latter being preferably used in the presence of a suitable catalyst such as, for example, palladium-on-charcoal, platinum-on-charcoal and the like. In order to prevent the undesired further hydrogenation of certain functional groups in the reactants and the reaction products it may be advantageous to add an appropriate catalyst-poison to the reaction mixture, e.g., thiophene and the like.

The compounds of formula (I) can also be converted into each other following art-known procedures of functional group transformation. Some examples of such procedures will be cited hereinafter.

The hydrogen atom of the amino function(s) of compounds of formula (I) may be substituted following art-known procedures such as, for example N-alkylation, N-acylation, reductive N-alkylation and the like procedures. For example alkylcarbonyl, arylcarbonyl and the like groups may be introduced by reacting the starting amine with an appropriate carboxylic acid or a derivative thereof such as, for example, an acid halide, acid anhydride and the like.


In all of the foregoing and in the following preparations, the reaction products may be isolated from the reaction mixture and, if desired, further purified according to methodologies generally known in the art.

The compounds of formula (I) have basic properties and, consequently, they may be converted to their therapeutically active non-toxic acid addition salt forms by treatment with appropriate acids, such as, for example, inorganic acids, such as hydrohalic acid, e.g., hydrochloric, hydrobromic and the like, and sulfuric acid, nitric acid, phosphoric acid and the like; or organic acids, such as, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 2-oxopropanoic, ethanedioic, propanedioic, butanedioic, (Z)-2-butenedioic, (E)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxybutanedioic, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids. Conversely the salt form can be converted by treatment with alkali into the free base form.

Some of the intermediates and starting materials in the foregoing preparations are known compounds while others are novel. They may be prepared according to art-known methodologies of preparing said known or similarly known compounds. Some procedures for preparing such intermediates will be described hereinafter in more detail.

The intermediates of formula (II) can conveniently be prepared following art-known procedures as described in, for example, U.S. Patent Nos. 4,219,559, 4,556,660 and 4,588,722 which are incorporated herein by reference.

For example, by cyclodesulfurization of an intermediate of formula (XXIV) following the same procedures as described hereinabove for the preparation of (I) starting from (XIX) and, subsequently eliminating the protective group P in the thus obtained intermediate of formula (XXV).



In (XXIV) and (XXV) P represents a protective group which is readily removeable by hydrogenation or hydrolysis, such as, phenylmethyl, C_1-4 alkyloxycarbonyl, aryl C_1-4 alkyloxycarbonyl and the like group.

The intermediates of formula (II) can also be prepared by reacting a piperidine derivative of formula (XXVI) with a benzimidazole derivative of formula (XXIII), following the same procedures as described hereinabove for the preparation of (I) starting from (XXII) and (XXIII) and, subsequently eliminating the protective group P in the thus obtained intermediate of formula (XXV).



Intermediates of formulae (V), (VII), (IX) and (XI) can conveniently be prepared following art-known procedures as described in, for example, U.S. Patent No. 4,556,660 by N-alkylating an intermediate of formula (II) with an appropriate reagent following a similar N-alkylation procedure as described hereinabove for the preparation of (I) starting from (II) and (III).

From formula (I) it is evident that the compounds of this invention may have several asymmetric carbon atoms in their structure. Each of these chiral centers may be present in a R- and S-configuration, this R- and S-notation being in correspondence with the rules described by R.S. Cahn, C. Ingold and V. Prelog in Angew. Chem., Int. Ed. Engl., 5, 385, 511 (1966).

Further, the compounds of this invention wherein L is C_3-6 alkenyl optionally substituted with Ar^3 may be present as E- and Z-forms, this E- and Z-notation being in correspondence with the rules also described in J. Org. Chem., 35, 2849-2867 (1979).

Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereoisomers may be separated by physical separation methods such as selective crystallization and chromatographic techniques, e.g., counter current distribution, and enantiomers may be separated from each other by the selective crystallization of their diastereomeric salts with optically active acids.

Pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically.

It is evident that the cis and trans diastereomeric racemates may be further resolved into their optical isomers, cis(+), cis(-), trans(+) and trans(-) by the application of methodologies known to those skilled in the art.

Stereochemically isomeric forms of the compounds of formula (I) are naturally intended to be embraced within the scope of the invention.

The compounds of formula (I), the pharmaceutically acceptable acid-addition salts and possible stereochemically isomeric forms thereof possess useful pharmacological properties. More particularly, they are active as anti-histaminics which activity is clearly evidenced by e.g. the results obtained in the "Protection of Rats from Compound 48/80-induced lethality"-test, the "Histamine antagonism in Guinea-Pig"-test and the "Ascaris Allergy test in Dogs"-test described in Arch. Int. Pharmacodyn. Ther. 251, 39-51 (1981). Apart from their anti-histaminic properties some of the subject compounds also show serotonin-antagonism.

Furthermore the compounds of formula (I), the pharmaceutically acceptable acid-addition salts and stereochemically isomeric forms thereof are particularly attractive due to their favourable pharmacokinetical profile. In particular, some compounds show a rapid onset so that their anti-histaminic effects are almost instantaneously present.

In view of their anti-histaminic properties, the compounds of formula (I) and their acid-addition salts are very useful in the treatment of allergic diseases such as, for example, allergic rhinitis, allergic conjunctivitis, chronic urticaria, allergic asthma and the like.

In view of their useful pharmacological properties the subject compounds may be formulated into various pharmaceutical forms for administration purposes. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compounds, in base or acid addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment. Acid addition salts of (I) due to their increased water solubility over the corresponding base form, are obviously more suitable in the preparation of aqueous compositions.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

The present invention is also related with a method of treating allergic diseases in warm-blooded animals suffering from said allergic diseases by administering an effective anti-allergic amount of a compound of formula (I) or a pharmaceutically acceptable acid addition salt thereof.

Those of skill in treating allergic diseases in warm-blooded animals could easily determine the effective amount from the test results presented hereinafter. In general it is contemplated that an effective amount would be from 0.001 mg/kg to 100 mg/kg body weight, and more preferably from 0.01 mg/kg to 1 mg/kg body weight.

The following examples are intended to illustrate and not to limit the scope of the present invention in all its aspects. Unless otherwise stated all parts therein are by weight.

EXPERIMENTAL PART

A. Preparation of Intermediates

Example 1

a) A mixture of 12.9 parts of 4-methyl-2-furancarboxaldehyde, 9.1 parts of hydroxylamine hydrochloride, 11.9 parts of pyridine and 160 parts of methanol was stirred overnight at room temperature. The reaction mixture was evaporated and the residue was taken up in water. The whole was acidified with concentrated hydrochloric acid and the product was extracted with 1,1'-oxybisethane. The extract was washed with water, dried, filtered and evaporated, yielding 14 parts (95.6%) of 4-methyl-2-furancarboxaldehyde,oxime as a residue (int. 1).

b) A mixture of 14 parts of 4-methyl-2-furancarboxaldehyde,oxime, 400 parts of methanol and 40 parts of 2-propanol, saturated with hydrogen chloride was hydrogenated at normal pressure and at room temperature with 4 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated at 30° C. The residue was taken up in water and saturated with potassium carbonate. The product was extracted with 1,1'-oxybisethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated, yielding 8 parts (64.2%) of 4-methyl-2-furanmethanamine as a residue (int. 2).

c) A mixture of 10.4 parts of 2-chloro-3-nitropyridine, 8 parts of 4-methyl-2-furanmethanamine, 7.6 parts of sodium hydrogen carbonate and 120 parts of ethanol was stirred overnight at reflux temperature. The reaction mixture was evaporated and the residue was taken up in water. The product was extracted with 1,1'-oxybisethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using trichloromethane as eluent. The pure fractions were collected and the eluent was evaporated, yielding 14.9 parts (91.2%) of N-[(4-methyl-2-furanyl)methyl]-3-nitro-2-pyridinamine as a residue (int. 3).

d) A mixture of 14.9 parts of N-[(4-methyl-2-furanyl)methyl]-3-nitro-2-pyridinamine, 2 parts of a solution of thiophene in methanol 4% and 320 parts of methanol was hydrogenated at normal pressure and at room temperature with 2 parts of platinum-on-charcoal catalyst 5%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated, yielding 14.2 parts (100%) of N²-[(4-methyl-2-furanyl)methyl]-2,3-pyridinediamine as a residue (int. 4).

In a similar manner there were also prepared:

N¹-[(5-methyl-2-furanyl)methyl]-1,2-benzenediamine as a residue (int. 5);
 N²-[(5-methyl-2-furanyl)methyl]-2,3-pyridinediamine as a residue (int. 6);
 N²-[(2-methyl-3-furanyl)methyl]-2,3-pyridinediamine as a residue (int. 7);
 N²-[(5-ethyl-2-furanyl)methyl]-2,3-pyridinediamine as a residue (int. 8);
 N⁴-[(5-methyl-2-furanyl)methyl]-3,4-pyridinediamine as a residue (int. 9);
 N²-[(3-methyl-2-furanyl)methyl]-2,3-pyridinediamine as a residue (int. 10);
 N³-[(5-methyl-2-furanyl)methyl]-3,4-pyridinediamine as a residue (int. 11);
 and N²-[(5-(1-methylethyl)-2-furanyl)methyl]-2,3-pyridinediamine as a residue (int. 12);

Example 2

- a) A mixture of 68.5 parts of ethyl 4-isothiocyanato-1-piperidinecarboxylate, 58.0 parts of N²-[(5-ethyl-2-furanyl)methyl]-2,3-pyridinediamine and 450 parts of tetrahydrofuran was stirred overnight at reflux temperature. After evaporation, the residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (97:3 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was stirred in 1,1'-oxybisethane. The product was filtered off and dried, yielding 50 parts (44.4%) of ethyl 4-[[[2-[(5-ethyl-2-furanyl)methyl]amino]-3-pyridinyl]amino]-thioxomethyl]amino]-1-piperidinecarboxylate (int. 13).
- b) A mixture of 50 parts of ethyl 4-[[[2-[(5-ethyl-2-furanyl)methyl]amino]-3-pyridinyl]amino]-thioxomethyl]amino]-1-piperidinecarboxylate, 32.4 parts of mercury(II) oxide and 480 parts of ethanol was stirred for 1 hour at reflux temperature. The reaction mixture was filtered over diatomaceous earth and the filtrate was evaporated. The residue was stirred in 1,1'-oxybisethane. The crystallized product was filtered off and dried, yielding 38 parts (83.1%) of ethyl 4-[[3-[(5-ethyl-2-furanyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinecarboxylate; mp. 111.1 °C (int. 14). In a similar manner there were also prepared:
- ethyl 4-[[1-[(5-methyl-2-furanyl)methyl]-1H-benzimidazol-2-yl]amino]-1-piperidinecarboxylate hemihydrate; mp. 150.1 °C (int. 15);
- ethyl 4-[[3-[(5-methyl-2-furanyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinecarboxylate as a residue (int. 16);
- ethyl 4-[[3-[(2-methyl-2-furanyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinecarboxylate; mp. 153.7 °C (int. 17);
- ethyl 4-[[1-[(5-methyl-2-furanyl)methyl]-1H-imidazo[4,5-c]pyridin-2-yl]amino]-1-piperidinecarboxylate; mp. 155.2 °C (int. 18);
- ethyl 4-[[3-[(3-methyl-2-furanyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinecarboxylate as a residue (int. 19);
- ethyl 4-[[3-[(5-methyl-2-furanyl)methyl]-3H-imidazo[4,5-c]pyridin-2-yl]amino]-1-piperidinecarboxylate (int. 20);
- ethyl 4-[[3-[(5-(1-methylethyl)-2-furanyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinecarboxylate as a residue (int. 21); and
- ethyl 4-[[3-[(4-methyl-2-furanyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinecarboxylate as a residue (int. 22);

Example 3

- To a stirred mixture of 4.6 parts of a sodium hydride dispersion 50% and 450 parts of N,N-dimethylformamide were added portionwise 57.6 parts of ethyl 4-(1H-benzimidazol-2-ylamino)-1-piperidinecarboxylate under nitrogen atmosphere. Upon complete addition, stirring was continued for 30 minutes at room temperature. 27.0 Parts of 3-(chloromethyl)-2-methylfuran were added dropwise while cooling. Upon completion, stirring was continued for 1 hour. Water was added dropwise to the mixture and the product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was stirred in 1,1'-oxybisethane. The solid product was filtered off and purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was stirred in 1,1'-oxybisethane. The product was filtered off and dried, yielding 37.5 parts (19.0%) of ethyl 4-[[1-[(2-methyl-3-furanyl)methyl]-1H-benzimidazol-2-yl]amino]-1-piperidinecarboxylate; mp. 150.4 °C (int. 23).

In a similar manner there was also prepared:

ethyl 4-[[1-4-thiazolylmethyl]-1H-benzimidazol-2-yl]amino]-1-piperidinecarboxylate; mp. 156.2 °C (int. 24)

Example 4

- A mixture of 20.5 parts of ethyl 4-[[1-[(5-methyl-2-furanyl)methyl]-1H-imidazo[4,5-c]pyridin-2-yl]amino]-1-piperidinecarboxylate, 40 parts of potassium hydroxide and 240 parts of 2-propanol was stirred overnight at reflux temperature. After evaporation, the residue was taken up in water and the product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was crystallized from 2-propanone. The product was filtered off and dried, yielding 15 parts (88%) of 1-[(5-methyl-2-furanyl)methyl]-N-(4-piperidinyl)-1H-imidazo[4,5-c]pyridin-2-amine; mp. 185.6 °C (int. 25).

In a similar manner there were also prepared:

- 1-[(5-methyl-2-furanyl)methyl]-N-(4-piperidiny)-1H-benzimidazol-2-amine as a residue (int. 26);
 N-(4-piperidiny)-1-(4-thiazolyl)methyl-1H-benzimidazol-2-amine dihydrobromide monohydrate; mp. 223.5 °C (int. 27);
- 5 3-[(5-methyl-2-furanyl)methyl]-N-(4-piperidiny)-3H-imidazo[4,5-b]pyridin-2-amine; mp. 120 °C (int. 28);
 3-[(2-methyl-3-furanyl)methyl]-N-(4-piperidiny)-3H-imidazo[4,5-b]pyridin-2-amine; mp. 165 °C (int. 29);
 3-[(5-methyl-2-furanyl)methyl]-N-(4-piperidiny)-3H-imidazo[4,5-b]pyridin-2-amine; mp. 106 °C (int. 30);
 1-[(2-methyl-3-furanyl)methyl]-N-(4-piperidiny)-1H-benzimidazol-2-amine; mp. 168 °C (int. 31);
 3-[(5-methyl-2-furanyl)methyl]-N-(4-piperidiny)-3H-imidazo[4,5-b]pyridin-2-amine; mp. 160 °C (int. 32);
- 10 3-[(5-ethyl-2-furanyl)methyl]-N-(4-piperidiny)-3H-imidazo[4,5-c]pyridin-2-amine hemihydrate; mp. 146 °C (int. 33);
 3-[(5-(1-methylethyl)-2-furanyl)methyl]-N-(4-piperidiny)-3H-imidazo[4,5-b]-pyridin-2-amine dihydrochloride; mp. 235 °C (int. 34); and
 3-[(4-methyl-2-furanyl)methyl]-N-(4-piperidiny)-3H-imidazo[4,5-b]pyridin-2-amine; mp. 143 °C (int. 35);

Example 5

- a) A mixture of 2 parts of 2-chloroacetonitrile, 8 parts of 1-[(5-methyl-2-furanyl)methyl]-N-(4-piperidiny)-1H-benzimidazol-2-amine, 3.1 parts of sodium carbonate and 90 parts of N,N-dimethylformamide was stirred and heated overnight at 45 °C. The reaction mixture was poured into water and the product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was crystallized from a mixture of acetonitrile and 2,2'-oxybispropane, yielding 7.3 parts (80.4%) of 4-[[1-[(5-methyl-2-furanyl)methyl]-1H-benzimidazol-2-yl]-amino]-1-piperidineacetonitrile; mp. 177.3 °C (int. 36).
- 25 b) A mixture of 6 parts of 4-[[1-[(5-methyl-2-furanyl)methyl]-1H-benzimidazol-2-yl]-amino]-1-piperidineacetonitrile and 200 parts of methanol, saturated with ammonia was hydrogenated at normal pressure and at room temperature with 2 parts of Raney-nickel catalyst. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated, yielding 6 parts of N-[1-(2-aminoethyl)-4-piperidiny]-1-[(5-methyl-2-furanyl)methyl]-1H-benzimidazol-2-amine as a residue (int. 37).

B. Preparation of Final Compounds

Example 6

- A mixture of 3.32 parts of 1-(2-bromoethyl)-4-ethyl-1,4-dihydro-5H-tetrazol-5-one, 4.65 parts of 1-[(5-methyl-2-furanyl)methyl]-N-(4-piperidiny)-1H-benzimidazol-2-amine, 1.6 parts of sodium carbonate and 45 parts of N,N-dimethylacetamide was stirred overnight at 80 °C. The reaction mixture was poured into water and the product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was purified by filtration over silica gel using a mixture of trichloromethane and methanol (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in methanol. The salt was filtered off and dried, yielding 5.2 parts (55%) of 1-ethyl-1,4-dihydro-4-[2-[[1-[(5-methyl-2-furanyl)methyl]-1H-benzimidazol-2-yl]-amino]-1-piperidiny]-ethyl]-5H-tetrazol-5-one ethanedioate(1:2); mp. 190.5 °C (compound 22).

Example 7

- A mixture of 3.9 parts of 2,3-dihydro-1,4-benzodioxin-2-methanol methanesulfonate(ester), 7.4 parts of 3-[(5-methyl-2-furanyl)methyl]-N-(4-piperidiny)-3H-imidazo[4,5-b]pyridin-2-amine, 5.3 parts of sodium carbonate and 90 parts of N,N-dimethylacetamide was stirred overnight at 80 °C. The mixture was poured into water and the product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of

trichloromethane and methanol (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in methanol. The salt was filtered off and dried, yielding 2.2 parts (23%) of N-[1-[(2,3-dihydro-1,4-benzodioxin-2-yl)methyl]-4-piperidinyl]-3-[5-methyl-2-furanyl)methyl]-3H-imidazo[4,5-b]pyridin-2-amine ethanedioate(1:2); mp. 222.3 °C (compound 14).

Example 8

A mixture of 1.2 parts of 1-chloro-2-ethoxyethane, 3.1 parts of 3-[(2-methyl-3-furanyl)methyl]-N-(4-piperidinyl)-3H-imidazo[4,5-b]pyridin-2-amine, 1.6 parts of sodium carbonate and 45 parts of N,N-dimethylformamide was stirred overnight at 70 °C. The reaction mixture was poured into 50 parts of water. Petroleum ether was added and after stirring, the crystallized product was filtered off, washed with water and petroleum ether and stirred in 1,1'-oxybisethane. The product was filtered off and dried at room temperature, yielding 1.6 parts (38.1%) of N-[1-(2-ethoxyethyl)-4-piperidinyl]-3-[(2-methyl-3-furanyl)methyl]-3H-imidazo[4,5-b]pyridin-2-amine dihydrate; mp. 84.5 °C (compound 32).

Example 9

A mixture of 1.82 parts of 3-bromo-1-propene, 7.4 parts of 3-[(5-methyl-2-furanyl)methyl]-N-(4-piperidinyl)-3H-imidazo[4,5-b]pyridin-2-amine ethanedioate (2:1), 4.2 parts of sodium hydrogen carbonate and 120 parts of ethanol was stirred overnight at reflux temperature. The reaction mixture was filtered and the filtrate was evaporated. The residue was taken up in water and the product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt in a mixture of 2-propanone and methanol. The salt was filtered off and dried, yielding 2 parts (23%) of 3-[(5-methyl-2-furanyl)methyl]-N-[1-(2-propenyl)-4-piperidinyl]-3H-imidazo[4,5-b]pyridin-2-amine (E)-2-butenedioate(1:2); mp. 172.7 °C (compound 5).

Example 10

A mixture of 2 parts of poly(oxyethylene), 4.5 parts of 3-[(5-methyl-2-furanyl)methyl]-N-(4-piperidinyl)-3H-imidazo[4,5-b]pyridin-2-amine, 1 part of a solution of thiophene in methanol 4% and 120 parts of methanol was hydrogenated at normal pressure and at 50 °C with 2 parts of platinum-on-charcoal catalyst 5%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off over diatomaceous earth and the filtrate was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile, yielding 0.7 parts (14.3%) of 3-[(5-methyl-2-furanyl)methyl]-N-(1-methyl-4-piperidinyl)-3H-imidazo[4,5-b]pyridin-2-amine; mp. 131.0 °C (compound 1).

Example 11

A mixture of 5 parts of a solution of acetaldehyde in tetrahydrofuran, 3.1 parts of 3-[(5-methyl-2-furanyl)methyl]-N-(4-piperidinyl)-3H-imidazo[4,5-b]pyridin-2-amine, 4.5 parts of tetrahydrofuran, 1 part of a solution of thiophene in methanol 4% and 200 parts of methanol was hydrogenated at normal pressure and at room temperature with 1 part of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated. The residue was converted into the (E)-2-butenedioate salt in ethanol. The salt was filtered off and dried, yielding 2.2 parts (38.5%) of N-(1-ethyl-4-piperidinyl)-3-[(5-methyl-2-furanyl)methyl]-3H-imidazo[4,5-b]pyridin-2-amine (E)-2-butenedioate(1:2); mp. 205.4 °C (compound 4).

Example 12

A mixture of 1.1 parts of 3-butene-2-one, 4.7 parts of 3-[(5-methyl-2-furanyl)methyl]-N-(4-piperidinyl)-3H-imidazo[4,5-b]pyridin-2-amine and 120 parts of ethanol was stirred for 3 hours at reflux temperature. After evaporation, the residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in methanol. The salt was filtered off and dried overnight in vacuo at 100 °C, yielding 1.8 parts (21.3%) of 4-[4-[[3-[(5-methyl-2-furanyl)-methyl]-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinyl]-2-butanone ethanedioate(1:2); mp. 164.1 °C. (compound 41).

Example 13

A mixture of 4.5 part of 2-(phenoxymethyl)oxirane, 4.65 parts of 1-[(5-methyl-2-furanyl)methyl]-N-(4-piperidinyl)-1H-benzimidazol-2-amine and 120 parts of methanol was stirred overnight at room temperature. The precipitated product was filtered off and dried, yielding 3.1 parts (44.8%) of 4-[[1-[(5-methyl-2-furanyl)-methyl]-1H-benzimidazol-2-yl]-amino]-α-(phenoxymethyl)-1-piperidineethanol; mp. 164.8 °C (compound 25).

Example 14


A mixture of 0.7 parts of isocyanatomethane, 2.5 parts of N-[1-(2-aminoethyl)-4-piperidinyl]-1-[(5-methyl-2-furanyl)methyl]-1H-benzimidazol-2-amine and 135 parts of tetrahydrofuran was stirred for 3 hours at room temperature. The precipitated product was filtered off and dried, yielding 1.2 parts (41.1%) of N-methyl-N-[2-[4-[[1-[(5-methyl-2-furanyl) methyl]-1H-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]urea; mp. 200.1 °C (compound 40).

Example 15

To a stirred mixture of 11.5 parts of N-(1-methyl-4-piperidinyl)-1H-benzimidazol-2-amine and 144 parts of N,N-dimethylformamide were added 5 parts of a sodium hydride dispersion 50%. After stirring for 1 hour at room temperature, a solution of 12.8 parts of 2-(chloromethyl)pyrazine in N,N-dimethylformamide was added dropwise to the thus obtained mixture. Upon complete addition, stirring was continued for 2 hours at 50 °C. The reaction mixture was poured into water and the product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 1.5 parts (9.3%) of N-(1-methyl-4-piperidinyl)-1-(2-pyrazinylmethyl)-1H-benzimidazol-2-amine; mp. 169.3 °C (compound 69).

All other compounds listed in table 1 were obtained by analogous methods of preparation as described in examples 6-15, the actual method of preparation being indicated in column 2 ("Ex. no.").

Table 1



Comp. No.	Ex. No.	L	R ²	Alk-R ¹	$\begin{matrix} 1 & 2 & 3 & 4 \\ A & A & A & A \end{matrix}$	salt/base	mp(°C)
1	10	CH ₃ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	base	131.0
2	6	4-CH ₃ -O-C ₆ H ₄ -CH ₂ -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	(E)-2-butenedioate(2:3)	199.8
3	10	(CH ₃) ₂ -CH-	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	(E)-2-butenedioate(1:2)	164.4
4	11	CH ₃ -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	(E)-2-butenedioate(1:2)	205.4
5	9	CH ₂ =CH-CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	(E)-2-butenedioate(1:2)	172.7
6	6	C ₆ H ₅ -CH=CH-CH ₂ - E form	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	2 (COOH) ₂	203.6
7	6	CH ₃ -CH ₂ -O-CH ₂ -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	2 (COOH) ₂	204.0
8	6	C ₆ H ₅ -O-(CH ₂) ₃ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	1/2 H ₂ O	96.5
9	6	HN-C(=O)-N-CH ₂ -CH ₂ - 	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	base	238.5
10	6	CH ₃ -CH ₂ -N-C(=O)-N-CH ₂ -CH ₂ - 	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	2 (COOH) ₂	194.1
11	11	CH ₃ -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	2 (COOH) ₂	211.7




5
10
15
20
25
30
35
40
45
50
55




Comp. No.	Ex. No.	L	R ²	Alk-R ¹	$\begin{matrix} 1 & 2 & 3 & 4 \\ A = A - A = A \end{matrix}$	salt/base	mp(°C)
12	7		H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	2 (COOH) ₂	224.8
13	6	4-CH ₃ -O-C ₆ H ₄ -CH ₂ -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	2 (COOH) ₂	224.8
14	7		H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	2 (COOH) ₂	222.3
15	7		H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	2 (COOH) ₂	213.3
16	10	CH ₃ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	(E)-2-butenedioate(2:3)	204.4
17	6	CH ₃ -CH ₂ -O-CH ₂ -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	(E)-2-butenedioate(2:3)	175.9
18	6	CH ₃ -C(=O)-(CH ₂) ₃ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	2 (COOH) ₂	190.1
19	6	C ₆ H ₅ -O-(CH ₂) ₃ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	1/2 H ₂ O	125.0
20	6		H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	base	211.7
21	6	C ₆ H ₅ -CH=CH-CH ₂ - E form	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	2 (COOH) ₂	225.5
22	6		H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	2 (COOH) ₂	190.5
23	7		H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	2 (COOH) ₂	216.2
24	6	CH ₃ -C(=O)-(CH ₂) ₃ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	2 (COOH) ₂	207.6
25	13	C ₆ H ₅ -O-CH ₂ -CH(OH)-CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	base	164.8

5
10
15
20
25
30
35
40
45
50
55

Comp. N.	Ex. No.	L	R ²	Alk-R ¹	$\Lambda^1 = \Lambda^2 = \Lambda^3 = \Lambda^4$	salt/base	mp(°C)
26	13	$C_6H_5-O-CH_2-CH(OH)-CH_2-$	H	$CH_2-5-CH_3-2-furanyl$	$-N=CH-CH=CH-$	base	147.2
27	6	$CH_3-CH_2-O-C(=O)-NH-(CH_2)_2-$	H	$CH_2-5-CH_3-2-furanyl$	$-CH=CH-CH=CH-$	base	124.3
28	9	$CH_2=CH-CH_2-$	H	$CH_2-5-CH_3-2-furanyl$	$-CH=CH-CH=CH-$	2 H ₂ O	82.5
29	10	CH_3-	H	$CH_2-5-C_2H_5-2-furanyl$	$-N=CH-CH=CH-$	base	135.1
30	8	$CH_3-CH_2-O-CH_2-CH_2-$	H	$CH_2-5-C_2H_5-2-furanyl$	$-N=CH-CH=CH-$	2 (COOH) ₂	178.6
31	8	$4-CH_3-O-C_6H_4-CH_2-CH_2-$	H	$CH_2-5-C_2H_5-2-furanyl$	$-N=CH-CH=CH-$	base	100.0
32	8	$CH_3-CH_2-O-CH_2-CH_2-$	H	$CH_2-2-CH_3-3-furanyl$	$-N=CH-CH=CH-$	2 H ₂ O	84.5
33	8	$4-CH_3-O-C_6H_4-CH_2-CH_2-$	H	$CH_2-2-CH_3-3-furanyl$	$-N=CH-CH=CH-$	base	117.5
34	6	$CH_3-CH_2-O-C(=O)-NH-(CH_2)_2-$	H	$CH_2-5-CH_3-2-furanyl$	$-N=CH-CH=CH-$	base	194.0
35	14	$CH_3-NH-C(=O)-NH-(CH_2)_2-$	H	$CH_2-5-CH_3-2-furanyl$	$-N=CH-CH=CH-$	2 HCl · H ₂ O	245.4
36	10	CH_3-	H	$CH_2-2-CH_3-3-furanyl$	$-N=CH-CH=CH-$	2 (COOH) ₂	202.0
37	6	$CH_3-(CH_2)_3-$	H	$CH_2-5-CH_3-2-furanyl$	$-N=CH-CH=CH-$	base	201.0
38	6		H	$CH_2-5-CH_3-2-furanyl$	$-N=CH-CH=CH-$	base	192.2
39	6	$CH_3-(CH_2)_5-$	H	$CH_2-5-CH_3-2-furanyl$	$-N=CH-CH=CH-$	2 (COOH) ₂	200.1
40	14	$CH_3-NH-C(=O)-NH-(CH_2)_2-$	H	$CH_2-5-CH_3-2-furanyl$	$-CH=CH-CH=CH-$	base	164.1
41	12	$CH_3-(C=O)-CH_2-CH_2-$	H	$CH_2-5-CH_3-2-furanyl$	$-N=CH-CH=CH-$	2 (COOH) ₂	188.5
42	6	$CH_3-(CH_2)_4-$	H	$CH_2-5-CH_3-2-furanyl$	$-N=CH-CH=CH-$	(E)-2-butenedioate(2:3)	179.9
43	10	CH_3-	H	$CH_2-5-CH_3-2-furanyl$	$-CH=CH-N=CH-$	base	96.1
44	6	$CH_3-CH_2-O-CH_2-CH_2-$	H	$CH_2-5-CH_3-2-furanyl$	$-CH=CH-N=CH-$	base	

5
10
15
20
25
30
35
40
45
50
55

Comp. No.	Ex. No.	L	R ²	Alk-R ¹	$\begin{matrix} 1 & 2 & 3 & 4 \\ A = A & - A & = A & \end{matrix}$	salt/base	mp(°C)
45	6	CH ₃ -CH ₂ -O-CH ₂ -CH ₂ -	H	CH ₂ -2-CH ₃ -3-furanyl	-CH=CH-CH=CH-	2 H ₂ O	110.7
46	6	4-F-C ₆ H ₄ -C(=O)-(CH ₂) ₃ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	H ₂ O	94.2
47	6	CH ₃ -C(=O)-CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	2 (COOH) ₂	184.2
48	6	4-F-C ₆ H ₄ -C(=O)-(CH ₂) ₃ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-CH=CH-	H ₂ O	109.2
49	6	4-CH ₃ -O-C ₆ H ₄ -CH ₂ -CH ₂ -	H	CH ₂ -2-CH ₃ -3-furanyl	-CH=CH-CH=CH-	base	164.0
50	9	CH ₂ =CH-CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=CH-N=CH-	base	165.8
51	10	CH ₃ -	H	CH ₂ -2-CH ₃ -3-furanyl	-CH=CH-CH=CH-	2 H ₂ O	102.3
52	6	4-CH ₃ -O-C ₆ H ₄ -CH ₂ -CH ₂ -	H	CH ₂ -3-CH ₃ -2-furanyl	-N=CH-CH=CH-	(E)-2-butenedioate(2:3)	218.8
53	6	 -CH ₂ -	H	CH ₂ -2-CH ₃ -3-furanyl	-N=CH-CH=CH-	base	212.8
54	6	 -CH ₂ -	H	CH ₂ -5-C ₂ H ₅ -2-furanyl	-N=CH-CH=CH-	1/2 H ₂ O	94.4
55	10	CH ₃ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=N-CH=CH-	2 (COOH) ₂	196.4
56	10	CH ₃ -	H	CH ₂ -3-CH ₃ -2-furanyl	-N=CH-CH=CH-	base	132.6
57	6	CH ₃ -CH ₂ -O-CH ₂ -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-CH=N-CH=CH-	2 (COOH) ₂	135.6
58	6	4-CH ₃ O-C ₆ H ₄ -CH ₂ -CH ₂ -	H	CH ₂ -5- <i>i</i> C ₃ H ₇ -2-furanyl	-N=CH-CH=CH-	2 HCl · H ₂ O	198.0
59	6	 -CH ₂ -	H	CH ₂ -5- <i>i</i> C ₃ H ₇ -2-furanyl	-N=CH-CH=CH-	base	180.3

Comp. No.	Ex. No.	L	R ²	Alk-R ¹	$\begin{matrix} 1 & 2 & 3 & 4 \\ A=A & -A & =A \end{matrix}$	salt/base	mp(°C)
60	8	CH ₃ -CH ₂ -O-CH ₂ -CH ₂ -	H	CH ₂ -5-IC ₃ H ₇ -2-furanyl	-N=CH-CH=CH-	(E)-2-butenedioate(1:2)	181.0
61	10	CH ₃ -	H	CH ₂ -5-IC ₃ H ₇ -2-furanyl	-N=CH-CH=CH-	2 HCl	257.1
62	10	CH ₃ -	H	CH ₂ -4-CH ₃ -2-furanyl	-N=CH-CH=CH-	(E)-2-butenedioate(1:2)	200.0
63	6	 -CH ₂ -	H	CH ₂ -4-CH ₃ -2-furanyl	-N=CH-CH=CH-	base	189.5
64	6	4-CH ₃ -O-C ₆ H ₄ -CH ₂ -CH ₂ -	H	CH ₂ -4-CH ₃ -2-furanyl	-N=CH-CH=CH-	(E)-2-butenedioate(1:2)	199.9
65	9	 -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	3 HCl	238.2
66	9	C ₆ H ₅ -S-CH ₂ -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	2 (COOH) ₂	200.8
67	11	CH ₃ -	H	CH ₂ -4-thiazolyl	-CH=CH-CH=CH-	2 HCl/H ₂ O	186.6
68	9	C ₆ H ₅ -SO ₂ -CH ₂ -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	2 (COOH) ₂	221.9
69	15	CH ₃ -	H	CH ₂ -2-pyrazinyl	-CH=CH-CH=CH-	base	169.3
70	9	 -CH ₂ -	H	CH ₂ -5-CH ₃ -2-furanyl	-N=CH-CH=CH-	3 HCl	232.5

C) Pharmacological Examples

The useful anti-histaminic properties of the compounds of formula (I) which can be used as the active ingredient in the formulations according to the present invention can be demonstrated by the following test procedure.

Example 16Protection of rats from compound 48/80-induced lethality.

Compound 48/80, a mixture of oligomers obtained by condensation of 4-methoxy-N-methylbenzeneethanamine and formaldehyde has been described as a potent histamine releasing agent (Int. Arch. Allergy, 13, 336 (1958)). The protection from compound 48/80-induced lethal circulatory collapse appears to be a simple way of evaluating quantitatively the antihistaminic activity of test compounds. Male rats of an inbred Wistar strain, weighing 240-260 g were used in the experiment. After overnight starvation the rats were transferred to conditioned laboratories (temp. = $21 \pm 1^\circ \text{C}$, relative humidity = $65 \pm 5\%$).

The rats were treated subcutaneously or orally with a test compound or with the solvent (NaCl solution, 0.9%). One hour after treatment there was injected intravenously compound 48-80, freshly dissolved in water, at a dose of 0.5 mg/kg (0.2 ml/100 g of body weight). In control experiments, wherein 250 solvent-treated animals were injected with the standard dose of compound 48/80, not more than 2.8% of the animals survived after 4 hours. Survival after 4 hours is therefore considered to be a safe criterion of a protective effect of drug administration. The ED_{50} -values of the compounds of formula (I) are listed in table 2. Said ED_{50} -values are the values in mg/kg body weight at which the tested compounds protect 50% of the tested animals against compound 48/80-induced lethality.

Table 2

No.	compound 48/80 lethality test in rats-ED ₅₀ in mg/kg body weight
1	0.005
4	0.0025
5	0.005
6	0.04
7	0.02
8	0.04
9	0.02
10	0.04
11	0.01
13	0.04
15	0.04
18	0.01
22	0.04
26	0.04
29	0.04
36	0.04
37	0.02
38	0.02
39	0.04
40	0.02
41	0.01
42	0.04
46	0.04
53	0.04
56	0.04
62	0.005
63	0.04
64	0.04

D) Composition Examples

The following formulations exemplify typical pharmaceutical compositions in dosage unit form suitable for systemic administration to animal and human subjects in accordance with the instant invention.

"Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I) or a pharmaceutically acceptable acid addition salt thereof.

Example 17 : ORAL DROPS

500 g of the A.I. was dissolved in 0.5 l of 2-hydroxypropanoic acid and 1.5 l of the polyethylene glycol at 60–80 °C. After cooling to 30–40 °C there were added 35 l of polyethylene glycol and the mixture was stirred well. Then there were added a solution of 1750 g of sodium saccharin in 2.5 l of purified water and while stirring there were added 2.5 l of cocoa flavor and polyethylene glycol q.s. to a volume of 50 l, providing an oral drop solution comprising 10 mg of the A.I. per ml. The resulting solution was filled into suitable containers.

Example 18 : ORAL SOLUTION

9 g of methyl 4-hydroxybenzoate and 1 g of propyl 4-hydroxybenzoate were dissolved in 4 l of boiling purified water. In 3 l of this solution were dissolved first 10 g of 2,3-dihydroxybutanedioic acid and thereafter 20 g of the A.I. The latter solution was combined with the remaining part of the former solution and 12 l 1,2,3-propane-triol and 3 l of sorbitol 70% solution were added thereto. 40 g of sodium saccharin were dissolved in 0.5 l of water and 2 ml of raspberry and 2 ml of gooseberry essence were added. The latter solution was combined with the former, water was added q.s. to a volume of 20 l providing an oral solution comprising 20 mg of the active ingredient per teaspoonful (5 ml). The resulting solution was filled in suitable containers.

Example 19 : CAPSULES

20 g of the A.I., 6 g sodium lauryl sulfate, 56 g starch, 56 g lactose, 0.8 g colloidal silicon dioxide, and 1.2 g magnesium stearate were vigorously stirred together. The resulting mixture was subsequently filled into 1000 suitable hardened gelating capsules, comprising each 20 mg of the active ingredient.

Example 20 : FILM-COATED TABLETSPreparation of tablet core

A mixture of 100 g of the A.I., 570 g lactose and 200 g starch was mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and 10 g polyvinylpyrrolidone (Kollidon-K-90®) in about 200 ml of water. The wet powder mixture was sieved, dried and sieved again. Then there was added 100 g microcrystalline cellulose (Avicel®) and 15 g hydrogenated vegetable oil (Sterotex®). The whole was mixed well and compressed into tablets, giving 10,000 tablets, each containing 10 mg of the active ingredient.

Coating

To a solution of 10 g methyl cellulose (Methocel 60 HG®) in 75 ml of denaturated ethanol there was added a solution of 5 g of ethyl cellulose (Ethocel 22 cps®) in 150 ml of dichloromethane. Then there were added 75 ml of dichloromethane and 2.5 ml 1,2,3-propane-triol. 10 g of polyethylene glycol was molten and

dissolved in 75 ml of dichloromethane. The latter solution was added to the former and then there were added 2.5 g of magnesium octadecanoate, 5 g of polyvinylpyrrolidone and 30 ml of concentrated colour suspension (Opaspray K-1-2109®) and the whole was homogenated. The tablet cores were coated with the thus obtained mixture in a coating apparatus.

Example 21 : INJECTABLE SOLUTION

1.8 g methyl 4-hydroxybenzoate and 0.2 g propyl 4-hydroxybenzoate were dissolved in about 0.5 l of boiling water for injection. After cooling to about 50 °C there were added while stirring 4 g lactic acid, 0.05 g propylene glycol and 4 g of the A.I..


The solution was cooled to room temperature and supplemented with water for injection q.s. ad 1 l volume, giving a solution of 4 mg A.I. per ml. The solution was sterilized by filtration (U.S.P. XVII p. 811) and filled in sterile containers.

Example 22 : SUPPOSITORIES

3 g A.I. was dissolved in a solution of 3 g 2,3-dihydroxybutanedioic acid in 25 ml polyethylene glycol 400. 12 g Surfactant (SPAN®) and triglycerides (Witepsol 555 ®) q.s. ad 300 g were molten together. The latter mixture was mixed well and the former solution. The thus obtained mixture was poured into moulds at a temperature of 37-38 °C to form 100 suppositories each containing 30 mg of the active ingredient.

Claims

1. A chemical compound having the formula



a pharmaceutically acceptable acid-addition salt or a stereochemically isomeric form thereof, wherein:

A¹ = A² = A³ = A⁴ is a bivalent radical having the formula

-CH=CH-CH=CH- (a-1),

-N=CH-CH=CH- (a-2),

-CH=N-CH=CH- (a-3),

-CH=CH-N=CH- (a-4), or

-CH=CH-CH=N- (a-5);

wherein one or two hydrogen atoms in said radicals (a-1)-(a-5) may, independently from each other, be replaced by halo, C₁₋₆alkyl, C₁₋₆alkyloxy, trifluoromethyl or hydroxy;

R is hydrogen or C₁₋₆alkyl;

R¹ is furanyl substituted with C₁₋₆alkyl, pyrazinyl, thiazolyl, or imidazolyl optionally substituted with C₁₋₆alkyl;

R² is hydrogen, C₁₋₆alkyl, C₃₋₆cycloalkyl, Ar³-C₁₋₆alkyl or (C₁₋₆alkyl)-CO;

L is C₃₋₆alkenyl optionally substituted with Ar³, or L is a radical of formula

-Alk-R³ (b-1),

-Alk-O-R⁴ (b-2),

-Alk-N(R⁵)-R⁶ (b-3),

-Alk-Z-C(=O)-R⁷ (b-4), or

-CH₂-CH(OH)-CH₂-O-R⁸ (b-5);

R³ is hydrogen, Ar¹-thio, Ar³-sulfonyl, 4,5-dihydro-5-oxo-1H-tetrazol-1-yl being optionally substituted in its 4-position with Ar³ or C₁₋₆alkyl, 2,3-dihydro-1,4-benzodioxin-2-yl-2,3-dihydro-1,4-benzodioxin-6-yl, 4-morpholinyl, 1-piperidinyl or 1-pyrrolidinyl, or when R¹ is furanyl substituted with C₁₋₆alkyl and A¹ = A² =

$A^2 = A^4$ is a bivalent radical of formula (a-1) or (a-2), R^3 may also be Ar^1 , 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl or 1-(C_1-5 alkyl)pyrrolyl;

R^4 is C_1-5 alkyl or Ar^1 ;

R^5 is C_1-5 alkyl optionally substituted with Ar^2 ;

5 R^6 is Ar^2 or C_1-5 alkyl optionally substituted with Ar^2 or when $A^1 = A^2 - A^3 = A^4$ is a radical of formula (a-1) or (a-2), R^6 may also be 1H-benzimidazol-2-yl;

R^7 is C_1-5 alkyl, Ar^2-C_1-5 alkyl, Ar^2 , amino, Ar^2 -amino, mono- and di(C_1-5 alkyl)amino, mono- and di(Ar^2-C_1-5 alkyl)amino, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl, C_3-5 cycloalkyloxy, C_1-5 alkyloxy or Ar^1-C_1-5 alkyloxy;

10 Z is O, NR^8 or a direct bond; said R^8 being hydrogen or C_1-5 alkyl optionally substituted with Ar^1 ;

R^9 is Ar^3 ;

each Alk independently being C_1-5 alkanediyl;

Ar^1 is phenyl, substituted phenyl, naphthalenyl, thienyl, halothienyl, C_1-5 alkyl substituted thienyl or furanyl, wherein said substituted phenyl is phenyl substituted with 1, 2 or 3 substituents, each independently selected from the group consisting of halo, hydroxy, nitro, cyano, trifluoromethyl, C_1-5 alkyl, 15 C_1-5 alkyloxy, C_1-5 alkylthio, mercapto, C_1-5 alkylsulfonyl, C_1-5 alkylsulfonyl(C_1-5 alkyl), phenyl- C_1-5 alkylsulfonyl, phenylsulfonyl(C_1-5 alkyl), amino, mono- and di(C_1-5 alkyl)amino, carboxy, C_1-5 alkyloxycarbonyl and C_1-5 alkylcarbonyl;

Ar^2 has the same meanings of Ar^1 , and may also be pyridinyl, mono- and di(C_1-5 alkyloxy)pyridinyl or furanyl substituted with C_1-5 alkyl; and

Ar^3 has the same meanings of Ar^2 , and may also be pyrazinyl, 1-(C_1-5 alkyl)pyrrolyl, thiazolyl or imidazolyl optionally substituted with C_1-5 alkyl;

provided that $A^1 = A^2 - A^3 = A^4$ is a radical of formula (a-1) or (a-2) when L is a radical of formula (b-4).

2. A chemical compound according to claim 1 wherein R is hydrogen, R^1 is furanyl substituted with 25 C_1-5 alkyl and R^2 is hydrogen or C_1-4 alkyl.

3. A chemical compound according to claim 2 wherein L is C_3-5 alkenyl optionally substituted with phenyl or substituted phenyl, or wherein L is a radical of formula (b-1), (b-2) or (b-5) wherein R^4 is C_1-5 alkyl, phenyl or substituted phenyl, R^7 is phenyl, substituted phenyl or C_1-5 alkyl optionally substituted with phenyl or substituted phenyl, R^8 is hydrogen C_1-5 alkyl, and R^9 is phenyl, substituted phenyl or 30 naphthalenyl.

4. A chemical compound according to claim 3 wherein L is a radical of formula (b-1) or (b-2) wherein R^3 is hydrogen, phenylsulfonyl, 4,5-dihydro-5-oxo-1H-tetrazol-1-yl being optionally substituted in its 4-position with C_1-4 alkyl, 2,3-dihydro-1,4-benzodioxin-2-yl, 4-morpholinyl, 1-piperidinyl or 1-pyrrolidinyl, or when R^1 is furanyl substituted with C_1-5 alkyl and $A^1 = A^2 - A^3 = A^4$ is a bivalent radical of formula (a-1) or (a-2), R^3 may 35 also be 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl, thienyl, furanyl, phenyl or substituted phenyl.

5. A chemical compound according to claim 1 wherein the compound is 3-[(5-methyl-2-furanyl)methyl]-N-(1-methyl-4-piperidinyl)-3H-imidazo[4,5-b]pyridin-2-amine.

6. A pharmaceutical composition comprising a suitable pharmaceutical carrier and as active ingredient a therapeutically effective amount of a compound as claimed in any one of claims 1 to 5.

40 7. An anti-histaminic pharmaceutical composition comprising a suitable pharmaceutical carrier and as active ingredient an effective anti-histaminic amount of a compound as claimed in any one of claims 1 to 5.

8. A method of preparing a pharmaceutical composition, characterized in that a therapeutically effective amount of a compound as defined in any of claims 1 to 5 is intimately mixed with suitable pharmaceutical carriers.

45 9. A compound as claimed in any one of claims 1 to 5 for use as a medicine.

10. A compound as claimed in any one of claims 1 to 5 for use as an anti-histaminic medicine.

11. A process for preparing a chemical compound as claimed in any one of claims 1 to 5, characterized by

50 a) N-alkylating a piperidine of formula H-D (II) with a reagent of formula L-W¹ (III), wherein W¹ represents a reactive leaving group, in a reaction-inert solvent;

b) reductively N-alkylating a piperidine of formula H-D (II) with a ketone or aldehyde of formula L' = O (IV), said L' = O being an intermediate of formula L'H₂ wherein two geminal hydrogen atoms are replaced by =O, in a reaction-inert solvent thus preparing a compound of formula L'H-D (I-a), wherein L' is a geminal bivalent radical comprising R^3-C_1-5 alkylidene, R^4-O-C_1-5 alkylidene, $R^6R^5-N-C_1-5$ alkylidene and 55 $R^7-C(=O)-Z-C_1-5$ alkylidene;

c) O-alkylating an intermediate of formula HO-Alk-D (V) with a reagent of formula $R^{4-a}-W^1$ (VI) wherein R^{4-a} is Ar^1 , in a reaction-inert solvent, thus preparing a compound of formula $R^{4-a}-O-Alk-D$ (I-b);

d) O-alkylating an intermediate of formula W'-Alk-D (VII) with a reagent of formula R^{4-a}-O-H (VIII) wherein R^{4-a} is Ar¹, in a reaction-inert solvent, thus preparing a compound of formula (I-b);

e) reacting an isocyanate of formula R^{7-a}-N=C=O (X) wherein R^{7-a}-NH- is amino, Ar²-amino, C₁₋₅-alkylamino or Ar²-C₁₋₅-alkylamino, with an intermediate of formula H-Z'-Alk-D (IX) wherein Z' is NR⁸ or O, in a reaction-inert solvent, thus preparing a compound of formula R^{7-a}-NH-C(=O)-Z'-Alk-D (I-c-1);


f) reacting a reagent of formula R^{7-b}-H (XII) wherein R^{7-b} has the previously defined meaning of R⁷ provided that it is other than Ar² or C₁₋₅-alkyl optionally substituted with Ar², with an isocyanate of formula O=C=N-Alk-D (XI) in a reaction-inert solvent, thus preparing a compound of formula R^{7-b}-C(=O)-NH-Alk-D (I-c-2);

g) reacting a reagent of formula R^{7-c}-C(=O)-OH (XIII), wherein R^{7-c} is Ar² or C₁₋₅-alkyl optionally substituted with Ar², with (IX), in a reaction-inert solvent, if desired, after converting the OH group in (XIII) in a reactive leaving group or by reacting (XIII) with (IX) in the presence of a reagent capable of forming esters or amides, thus preparing a compound of formula R^{7-c}-C(=O)-Z'-Alk-D (I-c-3);

h) reacting an alkenylene of formula L²-C₂₋₆-alkenediyl-H (XIV), wherein L² being Ar¹-thio, Ar³-sulfonyl, Ar², C₁₋₅-alkylcarbonyl, Ar²-C₁₋₅-alkylcarbonyl, Ar²-carbonyl, C₁₋₅-alkyloxycarbonyl, Ar¹-C₁₋₅-alkyloxycarbonyl or C₃₋₅-cycloalkyloxycarbonyl, with a piperidine of formula H-D (II), in a reaction-inert solvent, thus preparing a compound of formula L²-C₂₋₆-alkanediyl-D (I-d);

i) reacting a reagent of formula

20




25

with a piperidine of formula H-D (II), in a reaction-inert solvent, thus preparing a compound of formula R⁹-O-CH₂-CH(OH)-CH₂-D (I-e);

j) reducing an intermediate of formula C₁₋₅-alkyl-O-C(=O)-D (XVI) with a reductant in a reaction-inert solvent, thus preparing a compound of formula CH₃-D (I-f);

wherein D represents a radical of formula

30




35

wherein A¹ = A² = A³ = A⁴, R, R¹ and R² have the previously described meanings; or

40

k) N-alkylating an intermediate of formula

45




(XVII)

50


with a reagent of formula W-Alk-R¹ (XVIII) in a reaction-inert solvent, or

l) cyclodesulfurizing an intermediate of formula


55




10 which may in situ be prepared by condensing an isothiocyanate of formula




with a diamine of formula



25 with an alkylhalide, metal oxide or metal salt in a reaction-inert solvent; or
m) reacting a piperidine of formula



35 with a benzimidazole derivative of formula



45 in a reaction-inert solvent wherein in (XXII) and (XXIII) either

Q¹ is -NHR² and Q² is W, or

Q¹ is W¹ and Q² is -NHR², or

Q¹ is =O and Q² is -NHR²; or

50 optionally converting the compound of formula (I) into each other following art-known group transformation procedures, and, if desired, converting the compounds of formula (I) into a therapeutically active non-toxic acid-addition salt form by treatment with an appropriate acid or, conversely, converting the acid-addition salt into the free base form with alkali; and/or preparing stereochemically isomeric forms thereof.

55



DOCUMENTS CONSIDERED TO BE RELEVANT			EP 88201172.9
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
P,X	EP - A2 - 0 232 937 (JANSSEN) * Claims 1,2 * --	1,7	C 07 D 401/14 C 07 D 401/12 C 07 D 417/14
D,A	EP - B1 - 0 099 139 (JANSSEN) * Claims 1,5-7 * --	1,7	C 07 D 405/14 C 07 D 409/14 C 07 D 471/04
D,A	US - A - 4 219 559 (JANSSENS) * Abstract; claim 1 * ----	1,7	A 61 K 31/44 A 61 K 31/445 A 61 K 31/415 A 61 K 31/425 A 61 K 31/34
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			C 07 D 401/00 C 07 D 417/00 C 07 D 405/00 C 07 D 409/00 C 07 D 471/00
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 04-08-1988	Examiner HAMMER
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date O : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	